

Behandeling

Treatment schedule for injectable gonadotropin

When you have problems with your ovulation, you can get hormone injections to help the egg develop.

(073) 553 86 60

Monday to Friday, 8.30 – 12.00 and 13.30 – 16.30

What does the procedure look like?

Here you will find all information about the procedure

Injections

What does the treatment cycle look like?

When do you start treatment?

When your menstruation begins, you contact the centre of reproductive medicine.

You make an appointment for a 'start'-ultrasound. This ultrasound has to take place in the first 5 days of your cycle. A 'start'-ultrasound is only necessary if you used hormones the previous month.

How do you use the injections?

You start with the injections on the third day of your cycle. Your doctor will tell you what dosage you need to use and when to stop using the injections.

How to fill in the schedule?

Cycle day 1. The day your menstruation starts and persists. The colour of the blood will be bright red. If the menstruation starts after 18:00, the next day will be first cycle day.

You fill in the day of your menstruation above cycle day '1'. If you fill in the other dates, you will see when you need to come in for the first ultrasound.

(schedule example is in Dutch)

Datum																				
Cyclusdag	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	
Gonal-F	X	X																		
Echo										+										
Ovitrelle																				

Gonal-F injections

In a natural cycle, follicle stimulating hormone (FSH) causes follicles to grow in the ovaria. FSH is produced by the brain and released into the blood. FSH is administered through subcutaneous injections (underneath the skin). These injections are Gonal-F injections.

At cycle day 3 you start with the Gonal-F injections. You inject yourself between 16:00 - 20:00h each day. You continue with the injections until the doctor tells you to stop. At each ultrasound you will hear which dosage you need to use. Usually there is an ultrasound every 2-3 days, so we can determine the exact time you need to use the Ovitrelle injection.

Ovitrelle® pre-filled syringe

Human Chorionic Gonadotropin (HCG) is a hormone produced by the body during pregnancy. The doctor will prescribe Ovitrelle to trigger the ovulation (the release of eggs by the ovaries).

The doctor or nurse will tell you when to inject the Ovitrelle. You only need it once in your cycle.

Learning to inject

Both injections will be injected subcutaneous (underneath the skin). Your nurse at the centre of reproductive medicine will explain this to you. Your partner can learn to inject as well. You will see there is little that can go wrong. You are given the opportunity to make an appointment for instruction to learn to inject yourself.

What are possible side effects?

There are few known side effects of FSH and HCG. Sometimes swollen breasts, intestinal problems, headaches and mood swings can occur. You might notice some rash or sensitivity at the injection site.

Questions

If you have any questions, you can contact the nurses of the digital office (<https://vpgjbz.digitalepoli.nl/login>) of the centre of reproductive medicine.

In case of emergency (before of after office hours), please contact the emergency department of Gynaecology and Obstetrics at (073) 553 20 20.

Betrokken afdelingen

Voortplantingsgeneeskunde (<https://www.jeroenboschziekenhuis.nl/afdelingen/voortplantingsgeneeskunde>)